

REFINEMENTS AND SHARPNESS OF SOME INEQUALITIES
FOR MATHIEU TYPE SERIES

ŽIVORAD TOMOVSKI* AND DELČO LEŠKOVSKI**

Dedicated to Academician Blagoj Popov on the Occasion of His 85th Birthday

Abstract. In this paper the classical Qi's type inequalities [15] for Mathieu type series as well as for its alternating version [13] are studied. Some new refinements and sharpness of the inequalities given in [13, 15, 19, 20] will be presented.

1. INTRODUCTION AND PRELIMINARIES

The following familiar infinite series

$$S(r) = \sum_{n=1}^{\infty} \frac{2n}{(n^2 + r^2)^2} \quad (r \in \mathbb{R}) \quad (1.1)$$

is named after Emile Leonard Mathieu (1835 – 1890), who investigated it in his 1890 work [10] on elasticity of solid bodies. Bounds for this series are needed for the solution of boundary value problems for the biharmonic equations in a two-dimensional rectangular domain (see [16], p.258, eq. (54)). Inequality

$$S(r) < \frac{1}{r^2} \quad (1.2)$$

was conjectured in 1890 by Mathieu [10] and proved only in 1952 by L. Berg [2]. A very elegant and at the same time elementary proof of the inequalities

$$\frac{1}{r^2 + \frac{1}{2}} < S(r) < \frac{1}{r^2} \quad (1.3)$$

is given by E. Makai [11]. These inequalities are called Mathieus.

H.W. Gould and T. A. Chapman [7] have proved in 1962 the inequalities

$$\frac{1}{r^2 + 1} - \frac{1}{r^2 + (m + 1)^2} \leq \sum_{n=1}^m \frac{2n}{(n^2 + r^2)^2} \leq \frac{1}{r^2} - \frac{1}{r^2 + m^2} \quad (\forall m \in \mathbb{N}) \quad (1.4)$$

from which we can derive weaker inequalities than (1.3).

A remarkably useful integral representation for $S(r)$ in the elegant form

$$S(r) = \frac{1}{r} \int_0^{\infty} \frac{t \sin(rt)}{e^t - 1} dt \quad (1.5)$$

2000 *Mathematics Subject Classification.* 26D15, 33C05, 33C10, 44A10, 33E20, 40A30.

Key words and phrases. Mathieu's series, inequalities, integral representations, Laplace transform, generalized hypergeometric functions, Bessel function of first kind, Riemann-Zeta function, Apéry's constant, two-sided inequality, generalized Mathieu series, trapezoidal quadrature rule, incomplete gamma function, Beta function, Gautschi's inequality.

was given by Emersleben [5]. Alternative version of (1.1)

$$\tilde{S}(r) = \sum_{n=1}^{\infty} (-1)^{n-1} \frac{2n}{(n^2 + r^2)^2} \quad (r \in \mathbb{R}) \quad (1.6)$$

was recently introduced by Pogany et.al in [13].

Let F be a Laplace transform of f i.e.

$$F(p) = L(f(t)) \quad (1.7)$$

Using the relations (see [14], pp.651)

$$\sum_{n=1}^{\infty} F(n) = \int_0^{\infty} \frac{f(x)}{e^x - 1} dx \quad (1.8)$$

$$\sum_{n=1}^{\infty} (-1)^{n-1} F(n) = - \int_0^{\infty} \frac{f(x)}{e^x + 1} dx \quad (1.9)$$

we gave [13] an integral representation for $\tilde{S}(r)$:

$$\tilde{S}(r) = \frac{1}{r} \int_0^{\infty} \frac{t \sin(rt)}{e^t + 1} dt \quad (1.10)$$

Several interesting problems and solutions dealing with integral representations and bounds for the following slight generalization of the Mathieu series with a fractional power

$$S_{\mu}(r) = \sum_{n=1}^{\infty} \frac{2n}{(n^2 + r^2)^{\mu}} \quad (r \in \mathbb{R}; \mu > 1) \quad (1.11)$$

can be found in the works by Diananda [4], Tomovski and Trencovski [18] and Cerone and Lenard [3]. Feng Qi introduced [15] the series of following type:

$$S_{\mu+1}(r, \alpha) = \sum_{n=1}^{\infty} \frac{2n^{\frac{\alpha}{2}}}{(n^{\alpha} + r^2)^{\mu+1}} \quad (r > 0, \mu > 0, \alpha > 0) \quad (1.12)$$

It is obvious that $S_{\mu}(r, 2) = S_{\mu}(r)$. Motivated essentially by the works of Cerone and Lenard [3] (and Qi [15]), Srivastava and Tomovski defined in [17] a family of generalized Mathieu series

$$S_{\mu}^{(\alpha, \beta)}(r, a) = S_{\mu}^{(\alpha, \beta)}(r, \{a_k\}_{k=1}^{\infty}) = \sum_{n=1}^{\infty} \frac{2a_n^{\beta}}{(a_n^{\alpha} + r^2)^{\mu}} \quad (r, \alpha, \beta, \mu \in \mathbb{R}) \quad (1.13)$$

where it is tacitly assumed that the positive sequence

$$a = \{a_k\}_{k=1}^{\infty} = \{a_1, a_2, a_3, \dots\} \quad (\lim_{k \rightarrow \infty} a_k = \infty)$$

is chosen such that the infinite series in definition (1.13) converges, that is, that the following auxiliary series

$$\sum_{n=1}^{\infty} \frac{1}{a_n^{\mu\alpha - \beta}}$$

is convergent. Comparing the definitions (1.1), (1.11) and (1.13), we see that $S_2(r) = S(r)$ and $S_{\mu}(r) = S_{\mu}^{(2,1)}(r, \{k\})$. Furthermore, the special cases

$S_2^{(2,1)}(r; \{a_k\})$, $S_\mu^{(2,1)}(r; \{k\})$, $S_\mu^{(2,1)}(r; \{k^\gamma\})$, $S_\mu^{(\alpha, \frac{\alpha}{2})}(r; \{k\}) = S_\mu(r, \alpha)$ were investigated by Qi [15]; Diananda [3]; Tomovski [19] and Cerone-Lenard [3].

Let

$$\begin{aligned} \tilde{S}_\mu^{(\alpha, \beta)}(r; a) &= \tilde{S}_\mu^{(\alpha, \beta)}(r; \{a_k\}_{k=1}^\infty) \\ &= \sum_{n=1}^\infty (-1)^{n-1} \frac{2a_n^\beta}{(a_n^\alpha + r^2)^\mu} \quad (r, \alpha, \beta, \mu \in \mathbb{R}) \end{aligned} \tag{1.14}$$

be an alternating variant of (1.13), where the positive sequence $\{a_n\}_{n=1}^\infty$ satisfied the same conditions of the definition (1.13). In [13] we obtained several integral representations for (1.13) and its alternating variant in terms of the generalized hypergeometric functions and the Bessel function of first kind. Using these integral representations it is shown [20] that

$$\tilde{S}_\mu^{(2,1)}(r; \{n^\gamma\}) \leq \frac{2}{(1+r^2)^\mu} = M(r, \mu) \tag{1.15}$$

with $r, \gamma \in \mathbb{R}^+$, $\mu > \frac{1}{2}(1+r^2)$.

Specially for $\mu = 2, \gamma = 1$, the inequality (1.15) reduces to the inequality

$$\tilde{S}(r) \leq M(r, 2) \quad (0 < r < \sqrt{3}). \tag{1.16}$$

2. REFINEMENTS OF THE QI TYPE INEQUALITIES

In this part firstly we shall refine some inequalities of Qi type, given in [15] and [13]. We need the following proposition.

Proposition 2.1. For $x > 0$,

$$\frac{x}{2e^x} < \frac{x}{e^x + 1} < \frac{x}{e^x}. \tag{2.1}$$

The following Theorem appears erroneously (without the term $-3e^{-\pi/(2r)}$ on the right hand sided inequality) in the works by Feng Qi [15] and we will present here the correct formulation.

Theorem 2.1. For any positive number $r > 0$,

$$L \leq S(r) \leq R \tag{2.2}$$

where

$$\begin{aligned} L &= \frac{4(1+r^2)(e^{-\pi/r} + e^{-\pi/(2r)}) - 4r^2 - 1}{(e^{-\pi/r} - 1)(1+r^2)(1+4r^2)} \\ R &= \frac{(1+4r^2)(e^{-\pi/r} - e^{-\pi/(2r)}) - 4(1+r^2) - 3e^{-\pi/(2r)}}{(e^{-\pi/r} - 1)(1+r^2)(1+4r^2)}. \end{aligned}$$

Remark 2.1. This theorem was published six years after appearance of the very important paper by Alzer et al. [1]. Namely they proved the following theorem.

Theorem 2.2. For any positive number $r > 0$,

$$\frac{1}{k_1 + r^2} < S(r) < \frac{1}{k_2 + r^2} \tag{2.3}$$

where $k_1 = \frac{1}{2\zeta(3)}$ and $k_2 = \frac{1}{6}$ are the best possible constants. Here ζ denotes Riemann-Zeta function and $\zeta(3) \approx 1.20205$ is the Apéry's constant.

Remark 2.2. When $0 < r < 0,72149\dots$, the upper bound in (2.2) is better than that in (2.3)(see figure 1).

Qi in [15] remarked that when $0 < r < 2,9002\dots$, the lower bound in (2.2) is positive, and then is useful. But, it is not better than that in (2.3).




FIGURE 1

The proof of the next Theorem is similar to those of Theorem 2.1 in [15] and it is sufficient to be presented here one of them.

Theorem 2.3. For all $r > 0$,

$$\tilde{L} < \tilde{S}(r) < \tilde{R}, \quad (2.4)$$

where

$$\tilde{L} = \frac{2}{(1+r^2)^2} - \frac{1}{(1+r^2)^2(1-e^{-\pi/r})} - \frac{\pi e^{-\pi/r}}{2r(1+r^2)(1-e^{-\pi/r})^2}$$

and

$$\tilde{R} = \frac{1}{(1+r^2)^2} + \frac{1}{(1+r^2)^2(1-e^{-\pi/r})} + \frac{\pi e^{-\pi/r}}{2r(1+r^2)(1-e^{-\pi/r})^2}.$$

Proof. Since

$$\tilde{S}(r) = \frac{1}{r} \sum_{k=0}^{\infty} \left(\int_{2k\pi/r}^{(2k+1)\pi/r} \frac{x}{e^x+1} \sin(rx) dx + \int_{(2k+1)\pi/r}^{(2k+2)\pi/r} \frac{x}{e^x+1} \sin(rx) dx \right), \quad (2.5)$$

by applying the elementary two-sided inequality (2.1), we obtain

$$\begin{aligned} \sum_{k=0}^{\infty} \int_{2k\pi/r}^{(2k+1)\pi/r} \frac{x}{e^x+1} \sin(rx) dx &\leq \sum_{k=0}^{\infty} \int_{2k\pi/r}^{(2k+1)\pi/r} x e^{-x} \sin(rx) dx, \\ \sum_{k=0}^{\infty} \int_{(2k+1)\pi/r}^{(2k+2)\pi/r} \frac{x}{e^x+1} \sin(rx) dx &\leq \frac{1}{2} \sum_{k=0}^{\infty} \int_{(2k+1)\pi/r}^{(2k+2)\pi/r} x e^{-x} \sin(rx) dx. \end{aligned}$$

Applying the well-known integral formula (see [14], p.446)

$$\int_0^{\infty} x e^{-x} \sin(rx) dx = \frac{2r}{(1+r^2)^2}$$

we get

$$\begin{aligned} \tilde{S}(r) &\leq \frac{1}{r} \left(\sum_{k=0}^{\infty} \int_{2k\pi/r}^{(2k+1)\pi/r} x e^{-x} \sin(rx) dx + \frac{1}{2} \sum_{k=0}^{\infty} \int_{(2k+1)\pi/r}^{(2k+2)\pi/r} x e^{-x} \sin(rx) dx \right) \\ &= \frac{1}{r} \left(\frac{1}{2} \int_0^{\infty} x e^{-x} \sin(rx) dx + \frac{1}{2} \sum_{k=0}^{\infty} \int_{2k\pi/r}^{(2k+1)\pi/r} x e^{-x} \sin(rx) dx \right) \\ &= \frac{1}{(1+r^2)^2} + \frac{1}{2r} \sum_{k=0}^{\infty} \int_{2k\pi/r}^{(2k+1)\pi/r} x e^{-x} \sin(rx) dx = \tilde{R}. \end{aligned}$$

On the other hand,

$$\begin{aligned} \tilde{S}(r) &\geq \frac{1}{r} \left(\frac{1}{2} \sum_{k=0}^{\infty} \int_{2k\pi/r}^{(2k+1)\pi/r} x e^{-x} \sin(rx) dx + \sum_{k=0}^{\infty} \int_{(2k+1)\pi/r}^{(2k+2)\pi/r} x e^{-x} \sin(rx) dx \right) \\ &= \frac{1}{r} \left(\int_0^{\infty} x e^{-x} \sin(rx) dx - \frac{1}{2} \sum_{k=0}^{\infty} \int_{2k\pi/r}^{(2k+1)\pi/r} x e^{-x} \sin(rx) dx \right) \\ &= \frac{1}{(1+r^2)^2} - \frac{1}{2r} \sum_{k=0}^{\infty} \int_{2k\pi/r}^{(2k+1)\pi/r} x e^{-x} \sin(rx) dx = \tilde{L}. \end{aligned}$$

The proof is complete. \square

Remark 2.3. Since

$$\begin{aligned} \frac{1}{1 - e^{-\frac{\pi}{r}}} &> 1 \quad (\forall r > 0), \\ \frac{\pi e^{-\frac{\pi}{r}}}{2r(1+r^2)(1 - e^{-\frac{\pi}{r}})^2} &> 0 \quad (\forall r > 0) \end{aligned}$$

we obtain that

$$\tilde{R} > \frac{1}{(1+r^2)^2} + \frac{1}{(1+r^2)^2} = M(r, 2),$$

i.e. the inequality (1.16) for $0 < r < \sqrt{3}$ is stronger than the right hand sided inequality of (2.4).

Theorem 2.4. [13] For all $r > 0$,

$$\frac{4\zeta(3) - 3}{2(3r^2 + 2)(2\zeta(3)r^2 + 1)} < \tilde{S}(r) < \frac{12 - \zeta(3)}{2(6r^2 + 1)(\zeta(3)r^2 + 2)}. \quad (2.6)$$

Remark 2.4. It is easy to show that when $0 < r < 0,28668$, the upper bound in (1.16) is better than the upper bound in (2.6). Namely, if we denote

$$D(r) = M(r, 2) - \frac{12 - \zeta(3)}{2(6r^2 + 1)(\zeta(3)r^2 + 2)} = \frac{2}{(1 + r^2)^2} - \frac{12 - \zeta(3)}{2(6r^2 + 1)(\zeta(3)r^2 + 2)},$$

after simplifying, we have

$$D(r) = \frac{(25\zeta(3) - 12)r^4 + 6(\zeta(3) + 4)r^2 + (\zeta(3) - 4)}{2(1 + r^2)^2(6r^2 + 1)(\zeta(3)r^2 + 2)}.$$

Let $G(r) = (25\zeta(3) - 12)r^4 + 6(\zeta(3) + 4)r^2 + (\zeta(3) - 4)$. For $r > 0$, the function $G(r)$ is increasing and the positive root of the equation $G(r) = 0$ is

$$r_0 = \sqrt{\frac{-3(\zeta(3) + 4) + \sqrt{-16\zeta^2(3) + 184\zeta(3) + 94}}{25\zeta(3) - 12}} \approx 0,28668.$$

Therefore, $D(r) < 0 \Leftrightarrow G(r) < 0 \Leftrightarrow r \in (0, r_0)$.




FIGURE 2

When $0 < r < 0,912173\dots$, the lower bound in (2.4) is positive, and then it is useful. It is better than the lower bound in (2.6) when $0 < r < 0,808583\dots$ (see figure 2).

Applying the same technique as Qi did in [15], that is, by using the fact that $x^{-1}(e^x + 1)$ is an increasing function for $x > c_0 \approx 1,27846$ (c_0 is a root of the equation $e^x - xe^x + 1 = 0$), the well-known inequality $\frac{x}{e^x+1} < \frac{2}{e^{x/2}}$ would yield the following result, which appears erroneously in the works by Pogany et al. [13, Theorem 4] and we will present here the correct formulation.

Theorem 2.5. (a) For all $r \in \left(0, \frac{\pi}{c_0}\right)$,

$$\tilde{S}(r) \leq \frac{1}{r} \int_0^{\pi/r} \frac{x \sin(rx)}{e^x + 1} dx. \tag{2.7}$$

(b) For all $r > 0$,

$$\frac{1}{r} \int_0^{\pi/r} \frac{x \sin(rx)}{e^x + 1} dx \leq \frac{2 \left(1 + e^{-\pi/(2r)}\right)}{r^2 + \frac{1}{4}}. \tag{2.8}$$

3. INEQUALITIES FOR GENERALIZED MATHIEU SERIES

In this section we shall refine double inequalities given in [19] and prove that the inequality (1.15) is stronger than inequality of [13] given by

$$\tilde{S}_\mu^{(2,1)}(r; \{n^\gamma\}) \leq \frac{b_L \tilde{C}_\mu(r) \Gamma\left(\mu + \frac{1}{2}\right)}{\left(\mu - \frac{3}{2}\right)^{\frac{1}{3}}} = N_b(r, \mu) \quad \left(\mu > \frac{3}{2}\right) \tag{3.1}$$

where $b_L = 0,674885\dots$ and $\tilde{C}_\mu(r) = \frac{\sqrt{\pi}}{(2r)^{\mu - \frac{3}{2}} \Gamma(\mu)}$.

Theorem 3.1. [19] For $r > 0, \mu > 0, \alpha > 0$ the following integral representation for $S_{\mu+1}(r, \alpha)$ holds

$$S_{\mu+1}(r, \alpha) = \frac{2}{\Gamma(\mu + 1)} \int_0^\infty x^\mu e^{-r^2 x} g(x) dx, \tag{3.2}$$

where $g(x) = \sum_{n=1}^\infty n^{\alpha/2} e^{-n^\alpha x}$.

The following lemma can be found in [6], pp146-147.

Lemma 3.1. Let at the some $z > 0$ function f is increasing on $[0, z)$ and decreasing on $[z, \infty)$ and $\int_0^\infty f(x) dx$ converges. Then the inequalities

$$\int_0^\infty f(x) dx - \max_{x \in \mathbb{R}^+} f(x) < \sum_{n=1}^\infty f(n) < \int_0^\infty f(x) dx + \max_{x \in \mathbb{R}^+} f(x) - \int_0^1 f(x) dx \tag{3.3}$$

take place.

Theorem 3.2. If $r > 0, \mu > 0, \alpha > 0, \mu - \frac{1}{\alpha} > -\frac{1}{2}$ then

$$\begin{aligned} & \frac{2}{\alpha r^{2\mu - 2/\alpha + 1}} B\left(\frac{1}{\alpha} + \frac{1}{2}, \mu - \frac{1}{\alpha} + \frac{1}{2}\right) - \frac{2\Gamma\left(\mu + \frac{1}{2}\right)}{\sqrt{2e}\Gamma(\mu + 1)r^{2\mu + 1}} < S_{\mu+1}(r, \alpha) \\ & < \frac{2}{\left(\mu\alpha + \frac{\alpha}{2} - 1\right)(r^2 + 1)^{\mu + 1}} {}_2F_1\left(1, \mu + 1; \mu - \frac{1}{\alpha} + \frac{3}{2}; \frac{r^2}{1 + r^2}\right) \\ & \quad + \frac{2\Gamma\left(\mu + \frac{1}{2}\right)}{\sqrt{2e}\Gamma(\mu + 1)r^{2\mu + 1}}, \end{aligned} \tag{3.4}$$

where ${}_2F_1$ denotes the Gauss hypergeometric function and B is the Euler Beta function.

Proof. The proof of (3.4) follows directly by applying of Lemma 3.1 to $f(x) = x^{\alpha/2}e^{-x^\alpha}$, $x > 0$, $t > 0$. Since $\int_0^\infty f(x)dx = \frac{\Gamma(\frac{1}{\alpha} + \frac{1}{2})}{\alpha t^{1/\alpha + 1/2}}$ and $\max_{x \in \mathbb{R}^+} f(x) = \frac{1}{\sqrt{2et}}$, by (3.3) we obtain

$$\frac{\Gamma\left(\frac{1}{\alpha} + \frac{1}{2}\right)}{\alpha t^{1/\alpha + 1/2}} - \frac{1}{\sqrt{2et}} < g(t) < \frac{\Gamma\left(\frac{1}{\alpha} + \frac{1}{2}, t\right)}{\alpha t^{1/\alpha + 1/2}} + \frac{1}{\sqrt{2et}} \quad (3.5)$$

where $g(x)$ is the function defined in Theorem 3.1. and $\Gamma(a, x) = \int_x^\infty e^{-t}t^{a-1}dt$ is an incomplete gamma function. Hence by Theorem 3.1. we obtain

$$\begin{aligned} & \frac{2\Gamma\left(\frac{1}{\alpha} + \frac{1}{2}\right)}{\alpha\Gamma(\mu + 1)} \int_0^\infty t^{\mu-1/\alpha-1/2} e^{-r^2t} dt - \frac{2}{\sqrt{2e}\Gamma(\mu + 1)} \int_0^\infty t^{\mu-1/2} e^{-r^2t} dt < S_{\mu+1}(r, \alpha) \\ & < \frac{2}{\alpha\Gamma(\mu + 1)} \int_0^\infty \Gamma\left(\frac{1}{\alpha} + \frac{1}{2}, t\right) t^{\mu-1/\alpha-1/2} e^{-r^2t} dt + \frac{2}{\sqrt{2e}\Gamma(\mu + 1)} \int_0^\infty t^{\mu-1/2} e^{-r^2t} dt, \\ & \frac{2\Gamma\left(\frac{1}{\alpha} + \frac{1}{2}\right)\Gamma\left(\mu - \frac{1}{\alpha} + \frac{1}{2}\right)}{\alpha\Gamma(\mu + 1)r^{2\mu-2/\alpha+1}} - \frac{2\Gamma\left(\mu + \frac{1}{2}\right)}{\sqrt{2e}\Gamma(\mu + 1)r^{2\mu+1}} < S_{\mu+1}(r, \alpha) \\ & < \frac{2}{\alpha\Gamma(\mu + 1)} \int_0^\infty \Gamma\left(\frac{1}{\alpha} + \frac{1}{2}, t\right) t^{\mu-1/\alpha-1/2} e^{-r^2t} dt + \frac{2\Gamma\left(\mu + \frac{1}{2}\right)}{\sqrt{2e}\Gamma(\mu + 1)r^{2\mu+1}}. \end{aligned}$$

Applying the integral formula (see [8], page 663)

$$\begin{aligned} & \int_0^\infty \Gamma\left(\frac{1}{\alpha} + \frac{1}{2}, t\right) t^{\mu-\frac{1}{\alpha}-\frac{1}{2}} e^{-r^2t} dt \\ & = \frac{\Gamma(\mu + 1)}{\left(\mu - \frac{1}{\alpha} + \frac{1}{2}\right)(r^2 + 1)^{\mu+1}} {}_2F_1\left(1, \mu + 1; \mu - \frac{1}{\alpha} + \frac{3}{2}; \frac{r^2}{1 + r^2}\right) \\ & \quad (r > 0, \mu > 0, \alpha > 0) \end{aligned} \quad (3.6)$$

and definition of the Beta function

$$B\left(\frac{1}{\alpha} + \frac{1}{2}, \mu - \frac{1}{\alpha} + \frac{1}{2}\right) = \frac{\Gamma\left(\frac{1}{\alpha} + \frac{1}{2}\right)\Gamma\left(\mu - \frac{1}{\alpha} + \frac{1}{2}\right)}{\Gamma(\mu + 1)} \quad (3.7)$$

we finally get (3.3). \square

Specially for $\alpha = 2$ we get two sided inequality for $S_{\mu+1}(r)$.

Corollary 3.1. For $r, \mu \in \mathbb{R}^+$,

$$\frac{1}{\mu r^{2\mu}} - \frac{2\Gamma\left(\mu + \frac{1}{2}\right)}{\sqrt{2e}\Gamma(\mu + 1)r^{2\mu+1}} < S_{\mu+1}(r) < \frac{1}{(\mu\Gamma(\mu + 1))(r^2 + 1)^\mu} + \frac{2\Gamma\left(\mu + \frac{1}{2}\right)}{\sqrt{2e}\Gamma(\mu + 1)r^{2\mu+1}}. \quad (3.8)$$

The next result was proved recently by Tomovski and Hilfer in [20], but we shall give here a modified proof illustrated graphically.

Theorem 3.3. For $r \in \mathbb{R}^+$, $\mu > \frac{3}{2}$ the inequality (1.15) is stronger than inequality (3.1).

Proof. Let

$$P(r, \mu) = \frac{M(r, \mu)}{N_b(r, \mu)} = \frac{(\mu - 3/2)^{1/3}}{b_L \sqrt{2\pi}} \cdot \frac{\Gamma(\mu)}{\Gamma(\mu + 1/2)} \cdot \frac{2^\mu r^{\mu-3/2}}{(1+r^2)^\mu}. \quad (3.9)$$

We shall prove that $P(r, \mu) < 1$ for $\mu > \frac{3}{2}$ and $\forall r > 0$.

For the proof of this theorem we need Gautschi's inequality (see [9],[12, page 286])

$$\frac{\Gamma(\mu + \beta)}{\Gamma(\mu)} \geq \left(\mu - \frac{1-\beta}{2}\right)^\beta \quad \left(\mu > \frac{1-\beta}{2}, \beta \in [0, 1]\right) \quad (3.10)$$

which for $\beta = \frac{1}{2}$ gives

$$\frac{\Gamma(\mu)}{\Gamma(\mu + 1/2)} \leq \frac{1}{\sqrt{\mu - 1/4}} \quad \left(\mu > \frac{1}{4}\right) \quad (3.11)$$

The function $f(r) = \frac{r^{\mu-3/2}}{(1+r^2)^\mu}$ has a maximum at the point $r_0 = \sqrt{\frac{\mu-3/2}{\mu+3/2}}$.

Then,

$$\begin{aligned} P(r, \mu) &\leq P(r_0, \mu) = \frac{(\mu - 3/2)^{1/3}}{b_L \sqrt{2\pi}} \cdot \frac{\Gamma(\mu)}{\Gamma(\mu + 1/2)} \cdot \frac{2^\mu \left(\sqrt{\frac{\mu-3/2}{\mu+3/2}}\right)^{\mu-3/2}}{\left(1 + \left(\sqrt{\frac{\mu-3/2}{\mu+3/2}}\right)^2\right)^\mu} \\ &= \frac{(\mu - 3/2)^{1/3}}{b_L \sqrt{2\pi}} \cdot \frac{\Gamma(\mu)}{\Gamma(\mu + 1/2)} \cdot \frac{2^\mu \frac{(\mu-3/2)^{\mu/2-3/4}}{(\mu+3/2)^{\mu/2-3/4}}}{\frac{2^\mu \mu^\mu}{(\mu+3/2)^\mu}} \\ &\leq \frac{1}{b_L \sqrt{2\pi}} \cdot \frac{(\mu - 3/2)^{\mu/2-3/4} \cdot (\mu + 3/2)^{\mu/2+3/4}}{\mu^\mu} \cdot \frac{(\mu - 3/2)^{1/3}}{(\mu - 1/4)^{1/2}} \end{aligned} \quad (3.12)$$

$P(r_0, \mu) \leq 1$ for $\mu > \frac{3}{2}$ (see figure 3)




FIGURE 3

i) $\mu \geq 2$.

The function $f(x) = \frac{(x-3/2)^{1/3}}{(x-1/4)^{1/2}}$ has maximum at $x = 4$.
Therefore, since (3.12), we have

$$P(r, \mu) \leq \frac{1}{b_L \sqrt{2\pi}} \cdot \frac{\left(\frac{5}{2}\right)^{1/3}}{\left(\frac{15}{4}\right)^{1/2}} \cdot L(\mu) \quad (3.13)$$

where

$$L(\mu) = \frac{(\mu - 3/2)^{\mu/2-3/4} \cdot (\mu + 3/2)^{\mu/2+3/4}}{\mu^\mu}.$$

Let $a = \mu - 3/2$. It is clear that $a \geq 1/2$. Then

$$\begin{aligned} L(\mu) &= L(a + 3/2) = \frac{a^{a/2} \cdot (a + 3)^{(a+3)/2}}{\left(\frac{2a+3}{2}\right)^{\frac{2a+3}{2}}} \\ &= \sqrt{\frac{\left(1 + \frac{3}{a}\right)^{a+3}}{\left(1 + \frac{3}{2a}\right)^{2a+3}}} = \sqrt{\left(\frac{1 + \frac{3}{a}}{1 + \frac{3}{a} + \frac{9}{4a^2}}\right)^a} \cdot \sqrt{\left(\frac{1 + \frac{3}{a}}{1 + \frac{3}{2a}}\right)^3} \\ &\leq \sqrt{\left(\frac{1 + \frac{3}{a}}{1 + \frac{3}{2a}}\right)^3} = \sqrt{\left(1 + \frac{3}{2a+3}\right)^3} \leq \left(\frac{7}{4}\right)^{3/2} \end{aligned} \quad (3.14)$$

Since (3.13) and (3.14) we have

$$P(r, \mu) \leq \frac{1}{b_L \sqrt{2\pi}} \cdot \frac{\left(\frac{5}{2}\right)^{1/3}}{\left(\frac{15}{4}\right)^{1/2}} \cdot \left(\frac{7}{4}\right)^{3/2} \leq 0,59113 \cdot \frac{1,35721}{1,93649} \cdot 2,31504 \leq 0,96 < 1$$

So, for $\mu \geq 2$, it holds $P(r, \mu) < 1$.

ii) $\frac{3}{2} < \mu \leq 2$ ($\frac{9}{6} < \mu \leq \frac{12}{6}$).

We will rewrite (3.12) in the following form

$$P(r, \mu) \leq \frac{1}{b_L \sqrt{2\pi}} \cdot \frac{(\mu - 3/2)^{\mu/2-5/12} \cdot (\mu + 3/2)^{\mu/2+3/4}}{\mu^\mu \cdot (\mu - 1/4)^{1/2}}. \quad (3.15)$$

Since $0 < \mu - \frac{3}{2} < 1$ by (3.15), we have

a) ($\frac{3}{2} < \mu \leq \frac{5}{3}$)

$$P(r, \mu) \leq \frac{1}{b_L \sqrt{2\pi}} \cdot \frac{\left(\frac{1}{6}\right)^{1/3} \cdot \left(\frac{19}{6}\right)^{19/12}}{\left(\frac{3}{2}\right)^{3/2} \cdot \left(\frac{5}{4}\right)^{1/2}} < 0,9825 < 1.$$

b) ($\frac{5}{3} < \mu \leq \frac{11}{6}$)

$$P(r, \mu) \leq \frac{1}{b_L \sqrt{2\pi}} \cdot \frac{\left(\frac{1}{3}\right)^{5/12} \cdot \left(\frac{20}{6}\right)^{20/12}}{\left(\frac{5}{3}\right)^{5/3} \cdot \left(\frac{17}{12}\right)^{1/2}} < 0,99764 < 1.$$

c) ($\frac{11}{6} < \mu \leq 2$)

$$P(r, \mu) \leq \frac{1}{b_L \sqrt{2\pi}} \cdot \frac{\left(\frac{1}{2}\right)^{1/2} \cdot \left(\frac{7}{2}\right)^{7/4}}{\left(\frac{11}{6}\right)^{11/6} \cdot \left(\frac{19}{12}\right)^{1/2}} < 0,97931 < 1.$$

This completes the proof. \square

REFERENCES

- [1] H. Alzer, J. L. Brenner, O. G. Ruehr, On Mathieu's inequality, *J. Math. Anal. Appl.*, **218** (1998), 607-610.
- [2] L. Berg, Über eine Abschätzung von Mathieu, *Math. Nachr.*, **7** (1952), 257-259.
- [3] P. Cerone, C. T. Lenard, On integral forms of generalized Mathieu series, *J. Inequal. Pure Appl. Math.*, **4** (5) (2003), Art.100, 1-11 (electronic).
- [4] P. H. Diananda, Some inequalities related to an inequality of Mathieu, *Math. Ann.*, **250** (1980), 95-98.
- [5] O. Emersleben, Über die Reihe $\sum_{k=1}^{\infty} k/(k^2 + c^2)^2$, *Math. Ann.*, **125** (1952), 165-171.
- [6] N. Lain Fernandez, *Polynomial bases on the sphere*, Univ. Lübeck, Inst. für Mathematik (Dissertation), **162p** (2003)
- [7] H. W. Gould, T. A. Chapman, Some curious limits involving definite integrals, *Amer. Math. Monthly*, **69** (1962), 651-653.
- [8] I. S. Gradshteyn, I. M. Ryzhik, Table of Integrals, Series, and Products, *Academic Press*, New York and London, (1965).
- [9] I. Lazarevic, A. Lupas, Functional equations for Wallis and Gamma functions, *Univ. Beograd Publ. Elektrotehn. Fak. Ser. Mat. Fiz.*, **461-497** (1979), 245-251.
- [10] E. Mathieu, Traite de physique mathematique, Vol. VI-VII, part 2. Paris 1890.
- [11] E. Makai, On the inequality of Mathieu, *Publ. Math.*, Debrecen, **5** (1957), 204-205.
- [12] D. S. Mitrinovic, Analytic Inequalities, *Springer-Verlag*, Berlin-Heidelberg-New York, (1970).
- [13] T. K. Pogany, H. M. Srivastava, Ž. Tomovski, Some families of Mathieu **a**-series and alternating Mathieu **a**-series, *Appl. Math. Comput.*, **173** (2006), 69-108.
- [14] A. P. Prudnikov, Yu. A. Bryckov, O. I. Maricev, Integrals and Series (Elementary Functions), *Nauka*, Moscow, 1981 (Russian); English translation: Integrals and Series, Vol.1: Elementary Functions, *Gordon and Breach Science Publishers*, New York, 1990.
- [15] Feng Qi, An integral expression and some inequalities of Mathieu type series, *Rostock. Math.Kolloq.*, **58** (2004), 37-46.
- [16] K. Schroder, Das Problem der eingespannten rechteckigen elastischen Platte, *Math. Anal.*, **121** (1) (1949), 247-326.
- [17] H. M. Srivastava, Ž. Tomovski, Some problems and solutions involving Mathieus series and its generalizations, *J. Inequal. Pure Appl. Math.*, **5** (2) (2004), Article 45, 1-13 (electronic).
- [18] Ž. Tomovski, K. Trenčevski, On an open problem of Bai-Ni Guo and Feng Qi, *J. Inequal. Pure Appl. Math.*, **4** (2) (2003), Article 29, 1-7 (electronic).
- [19] Ž. Tomovski, New double inequalities for Mathieu type series, *Univ. Beograd Publ. Elektrotehn. Fak. Ser. Mat.*, **15** (2004), 79-83.
- [20] Ž. Tomovski, R. Hilfer, Some bounds for alternating Mathieu type series, *J. Math. Inequal.*, **2** (2008), 17-26.

*"ST. CYRIL AND METHODIUS UNIVERSITY"

FACULTY OF NATURAL SCIENCES AND MATHEMATICS
 INSTITUTE OF MATHEMATICS, P.O. BOX 162
 E-mail address: tomovski@iunona.pmf.ukim.edu.mk

**INTERNATIONAL BALKAN UNIVERSITY, SKOPJE

E-mail address: dleskovski@gmail.com